



# Ed Injections

---

COMPOUNDING WITHOUT  
COMPROMISE SINCE 1962

This drug was compounded for you by O'Brien Pharmacy according to your provider's prescription. **CAUTION** Federal law prohibits the transfer of this drug to any person other than the patient for whom it was prescribed.

This information is provided in addition to the written information and pharmacist consultation you received. No standardized patient information is available for this compound.

Prior to use, please read the detailed patient packet provided with this medication and keep it on hand for future reference.

## Use

Common uses include erectile dysfunction.

## Side Effects

It may cause headache, bruising at injection site, upset stomach, drowsiness, flushing, light head, or prolonged/painful erection called priapism.

## Precautions

Contact prescriber if redness, swelling, lumps, curving or priapism occurs. Seek medical attention if erection is painful or lasts longer than four hours. Do not use if solution is cloudy or has particles.

## Directions

Swab vial and skin with sterile alcohol before use. Use as directed by prescriber and do not exceed dose. Consult prescriber before using more than 3 times/week.

## Storage

Store in freezer until beyond use date indicated on the label. Warm to room temperature before filling syringe or injecting.

Do not dispose of in waterways. Do not flush unused medications or pour down a sink or drain. You have the right to know about the proper use of your medication and its effects. A pharmacist is available for consult Mon. - Fri. 10 a.m. to 6 p.m. Central Time by calling the pharmacy at 800-627-4360. Patients in California will receive a return call on Saturdays for voicemails left by noon Central Time (10 a.m. Pacific Time).